UNDERSTANDING ISOLATION OF AFFECT & HOW IT DEVELOPS IN MEDICAL STUDENTS

NEW YORK INSTITUTE OF TECHNOLOGY

College of Osteopathic Medicine _AUTHORS_

David Ozga MBS, OMS4, Imari Faceson OMS4, Alex Hernandez MS, OMS3, Colleen Kirk DPS, MBA, MIM
NYIT College of Osteopathic Medicine, Old Westbury, NY

NEW YORK INSTITUTE OF TECHNOLOGY

School of Management

INTRODUCTION

- Isolation of affect is the ability of an individual to mask their emotional state in an emotionally charged situation [1-3].
- Comparable to a physician delivering a terminal diagnosis to an emotional patient.
- Isolation of affect is a skill is taught to future physicians throughout medical school [1-5].
- Quantifying the degree of isolation and relevant factors has not been studied in this context [1-8].

Hypotheses:


- H1: Isolation of affect in a patient context will increase between 1st and 4th year medical students.
- H2: Trait crying proneness will decrease as medical students progress in their education.

OBJECTIVE

Through a mixed methods analysis, this research aims to investigate:

(1) Factors that impact the ability to isolate one's emotional state

(2) Key personality traits including communal narcissism, crying proneness, and empathy



METHODOLOGY

Mixed Methods

- Interviews with 4th-year medical students (N=5).
- Qualtrics survey administered via Amazon's Mechanical Turk platform (N=102).
- Virtual Interviews
- Qualtrics Surveys

— A two-tailed bivariate correlation analysis was performed between all variables using SPSS.

- Pearson coefficients are presented in this project.
- Qualitative interview data analyzed via theme coding and thematic mapping.

ANALYSIS





Fig. 1 Thematic Mapping of Medical Student Interview Data.

RESULTS/FINDINGS -


4th-year Medical Student Interview Quotes

- "My emotional reaction internally has lessened over time. I don't react as much to death and dying."
- "Being in the operating room is a particularly stressful experience, because your presence can either be a complete shadow or you could be yelled at just for existing there, especially if you're scrubbed in."
- "[Crying] is just a moment of vulnerability that in this day and age in medicine just isn't acceptable"
- "I would feel weak if other people witnessed me begin to cry."
- "I would feel others are judging me and thinking that I am not adept at being a clinician because I am unable to handle the situation."
- "I am more mild-mannered and less reactive than I was when I started medical school."



** Correlation is significant at the 0.01 level (2-tailed). *Correlation is significant at the 0.05 level (2-tailed). N=102 Fig. 2 Correlation Matrix







CONCLUSION

Future Research

- Compare 1st-year medical students to 4th-year medical students in their subjective level of isolation of affect.
- Survey medical students to quantify difference in affect between 1st and 4th year students.

